

ON A p -ADIC INTERPOLATION FUNCTION FOR THE q -EXTENSION OF THE GENERALIZED BERNOULLI POLYNOMIALS AND ITS DERIVATIVE

TAEKYUN KIM

*Institute of Science Education,
Kongju National University, Kongju 314-701, S. Korea
e-mail: tkim64@hanmail.net (or tkim@kongju.ac.kr)*

ABSTRACT. In [18], the new q -extension of Bernoulli polynomials and generalized Bernoulli numbers attached to χ were constructed by using p -adic invariant integral on $\mathbb{Z}_p$. In this paper we construct the new q -extension of generalized Bernoulli polynomials attached to χ due to author and derive the existence of a specific p -adic interpolation function which interpolate the q -extension of generalized Bernoulli polynomials at negative integer. Finally, we give the values of partial derivative for this function and investigate some properties which are related to this interpolation functions.

§1. INTRODUCTION

Let p be a fixed prime. Throughout this paper $\mathbb{Z}_p$, $\mathbb{Q}_p$, $\mathbb{C}$ and $\mathbb{C}_p$ will, respectively, denote the ring of p -adic rational integers, the field of p -adic rational numbers, the complex number field and the completion of algebraic closure of $\mathbb{Q}_p$, cf.[3, 10, 17]. Let v_p be the normalized exponential valuation of $\mathbb{C}_p$ with $|p|_p = p^{-v_p(p)} = p^{-1}$. When one talks of q -extension, q is variously considered as an indeterminate, a complex number $q \in \mathbb{C}$, or a p -adic number $q \in \mathbb{C}_p$. If $q \in \mathbb{C}$, one normally assumes $|q| < 1$. If $q \in \mathbb{C}_p$, then we assume $|q - 1|_p < p^{-\frac{1}{p-1}}$, so that $q^x = \exp(x \log q)$ for $|x|_p \leq 1$. Kubota and Leopoldt proved the existence of meromorphic functions, $L_p(s, \chi)$, defined over the

Key words and phrases. p -adic q -integrals, multiple Barnes' Bernoulli numbers.
2000 Mathematics Subject Classification: 11S80, 11B68, 11M99 .

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ -TEX

p -adic number field, that serve as p -adic equivalents of the Dirichlet L -series. These p -adic L -functions interpolate the values

$$L_p(1-n, \chi) = -\frac{1}{n}(1 - \chi_n(p)p^{n-1})B_{n, \chi_n}, \text{ for } n \in \mathbb{N} = \{1, 2, \dots\},$$

where $B_{n, \chi}$ denote the n th generalized Bernoulli numbers associated with the primitive Dirichlet character χ , and $\chi_n = \chi w^{-n}$, with w the *Teichmüller* character, cf.[1-24]. In a recent paper [18], the author constructed the new q -extensions of Riemann zeta function, Hurwitz's zeta function and Dirichlet's L -functions. In Section 2, we define the q -extension of generalized Bernoulli polynomials attached to χ and construct a new q -extension of Hurwitz's type L -function which interpolates the q -extension of generalized Bernoulli numbers attached to χ at negative integers. In $\mathbb{C}$, we introduce some of the basic facts about the q -extension of Hurwitz's type L -function which interpolates the generalized q -Bernoulli polynomials attached to χ . The values of this function at negative integers are algebraic, hence may be regarded as lying in an extension of $\mathbb{Q}_p$. We therefore look for a p -adic function which agrees with at negative integers. The purpose of this paper is to construct the new q -extension of generalized Bernoulli polynomials attached to χ due to author and derive the existence of a specific p -adic interpolation function which interpolate the q -extension of generalized Bernoulli polynomials at negative integer. Finally, we give the values of partial derivative for this p -adic function and investigate some properties which are related to this p -adic interpolation functions in Section 3.

2. AN q -EXTENSION OF DIRICHLET'S L -FUNCTION

In this section we assume that $q \in \mathbb{C}$ with $|q| < 1$ and $h \in \mathbb{Z}$. The classical Bernoulli polynomials are defined by

$$\frac{t}{e^t - 1} e^{xt} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!}, \quad |t| < 2\pi,$$

and the classical Bernoulli numbers are defined by $B_n = B_n(0)$. Let χ be a primitive Dirichlet character of conductor $f \in \mathbb{N}$. Then the generalized Bernoulli polynomials associated with χ , $B_{n, \chi}(x)$, are also defined by

$$(1) \quad \sum_{a=1}^f \frac{\chi(a) t e^{(a+x)t}}{e^{ft} - 1} = \sum_{n=0}^{\infty} \frac{B_{n, \chi}(x)}{n!} t^n, \text{ cf. [4, 18].}$$

Let us define the q -extension of Bernoulli polynomials as follows:

$$(2) \quad \frac{h \log q + t}{q^h e^t - 1} e^{xt} = \sum_{n=0}^{\infty} B_{n,q}^{(h)}(x) \frac{t^n}{n!}, \text{ cf. [4, 18] }.$$

We now consider the q -extension of the generalized Bernoulli polynomials attached to χ as

$$(3) \quad \frac{\sum_{i=0}^{f-1} (te^{it} \chi(i) q^{hi} + e^{ti} \log q^h q^{hi} \chi(i))}{q^{hf} e^{ft} - 1} e^{xt} = \sum_{n=0}^{\infty} B_{n,q,\chi}^{(h)}(x) \frac{t^n}{n!}, \quad |t| < 2\pi.$$

Let us define the q -extensions of Bernoulli numbers and generalized Bernoulli numbers attached to χ as $B_{n,q}^{(h)} = B_{n,q}^{(h)}(0)$ and $B_{n,q,\chi}^{(h)} = B_{n,q,\chi}^{(h)}(0)$. Then we note that

$$(4) \quad B_{n,q,\chi}^{(h)}(x) = \sum_{k=0}^n \binom{n}{k} B_{k,q,\chi}^{(h)} x^{n-k} = f^{n-1} \sum_{i=0}^{f-1} \chi(i) q^{hi} B_{n,q^f}^{(h)}\left(\frac{i+x}{f}\right).$$

Remark. Let $\delta_{1,k}$ be denoted by Kronecker symbol. Then we see that

$$B_{0,q}^{(h)} = \frac{h \log q}{q^h - 1}, \quad q^h (B_q^{(h)} + 1)^n - B_{n,q}^{(h)} = \delta_{1,k}, \quad k \geq 1, \text{ cf. [1, 2, 21, 22] },$$

with the usual convent of replacing $(B_q^{(h)})^n$ by $B_{n,q}^{(h)}$.

For $s \in \mathbb{C}$, the q -extension of Hurwitz zeta function is defined by

$$\zeta_q^{(h)}(s, x) = \sum_{n=0}^{\infty} \frac{q^{nh}}{(n+x)^s} - \frac{h \log q}{s-1} \sum_{n=0}^{\infty} \frac{q^{nh}}{(n+x)^{s-1}}, \text{ cf. [12, 14, 18].}$$

Note that $\zeta_q^{(h)}(s, x)$ is analytic continuation for $\Re(s) > 1$. This function have the below value at negative integers:

$$\zeta_q^{(h)}(1-n, x) = -\frac{B_{n,q}^{(h)}(x)}{n}, \text{ for } n \in \mathbb{N} = \{1, 2, \dots\}.$$

In [18], the q -extension of Dirichlet's L -function is also defined by

$$(5) \quad L_q^{(h)}(s, \chi) = \sum_{n=1}^{\infty} \frac{\chi(n) q^{hn}}{n^s} - \frac{h \log q}{s-1} \sum_{n=1}^{\infty} \frac{q^{hn} \chi(n)}{n^{s-1}}.$$

This function is analytic continuation for $\Re(s) > 1$. Note that $L_q^{(h)}(1-n, \chi) = -\frac{B_{n,q,\chi}^{(h)}}{n}$, for $n \in \mathbb{N}$, cf.[12, 14, 18, 19]. We now set

$$(6) \quad F_{q,\chi}^{(h)}(t, x) = \frac{\sum_{i=0}^{f-1} (te^{it}\chi(i)q^{hi} + e^{ti}\log q^h q^{hi}\chi(i))}{q^{hf}e^{ft} - 1} e^{xt}, \text{ for } |t| < \frac{2\pi}{f}.$$

By using (6), we easily see that

$$(7) \quad F_{q,\chi}^{(h)}(t, x) = -t \sum_{n=0}^{\infty} \chi(n) q^{nh} e^{(n+x)t} - h \log q \sum_{n=0}^{\infty} \chi(n) q^{hn} e^{(n+x)t}.$$

It is easy to see that the series on the right-hand side of (7) are uniformly convergent. Hence, we have

$$(8) \quad B_{k,q,\chi}^{(h)}(x) = \frac{d^k}{dt^k} F_{q,\chi}^{(h)}(t, x)|_{t=0} = -k \sum_{n=0}^{\infty} \chi(n) q^{hn} (n+x)^{k-1} - h \log q \sum_{n=0}^{\infty} \chi(n) q^{hn} (n+x)^k.$$

That is,

$$(9) \quad -\frac{B_{k,q,\chi}^{(h)}(x)}{k} = \sum_{n=0}^{\infty} \chi(n) q^{hn} (n+x)^{k-1} + \frac{h \log q}{k} \sum_{n=0}^{\infty} \chi(n) q^{hn} (n+x)^k, \quad k \in \mathbb{N}.$$

Thus, we can consider the q -extension of Dirichlet's L -function which interpolates the generalizes q -Bernoulli numbers at negative integer as follows:

Definition 1. For $s \in \mathbb{C}$, define

$$(10) \quad L_q^{(h)}(s, x|\chi) = \sum_{n=0}^{\infty} \frac{\chi(n) q^{hn}}{(n+x)^s} - \frac{h \log q}{s-1} \sum_{n=0}^{\infty} \frac{q^{hn} \chi(n)}{(n+x)^{s-1}}.$$

Note that $L_q^{(h)}(s, x|\chi)$ is analytic continuation in $\mathbb{C}$ with only simple pole at $s = 1$.

By (9) and (10), we obtain the following:

Proposition 2. For any positive integer k , we have

$$L_q^{(h)}(1-k, x|\chi) = -\frac{B_{k,q,\chi}^{(h)}}{k}(x).$$

We now give the integral representation of the q -extension of Dirichlet's L -function which interpolates the q -extension of generalized Bernoulli polynomials attached to χ in $\mathbb{C}$. Let $\Gamma(s)$ be the gamma function. Then we can readily see that

$$(11) \quad L_q^{(h)}(s, x|\chi) = \frac{1}{\Gamma(s)} \int_0^\infty t^{s-2} F_{q,\chi}^{(h)}(-t, x) dt.$$

From (3),(6) and (11), we note that

$$L_q^{(h)}(1-n, x|\chi) = -\frac{B_{n,q,\chi}^{(h)}(x)}{n}, \text{ for } n \in \mathbb{N}.$$

Let s be a complex variable, a and f be integers with $0 < a < f$. Then we define $H_q^{(h)}(s, a|f)$ as follows:

$$(12) \quad H_q^{(h)}(s, a|f) = \sum_{\substack{m \equiv a \pmod{f} \\ m > 0}} \frac{q^{mh}}{m^s} - \frac{\log q}{s-1} \sum_{\substack{m \equiv a \pmod{f} \\ m > 0}} \frac{q^{hm}}{m^{s-1}} = f^{-s} q^{ha} \zeta_{q^f}^{(h)}\left(s, \frac{a}{f}\right).$$

Let $\chi (\neq 1)$ be the Dirichlet's character with conductor $f \in \mathbb{N}$. Then the q -analogue of Dirichlet's L -function can be expressed as the sum.

$$(13) \quad L_q^{(h)}(s, \chi) = \sum_{a=1}^f \chi(a) H_q^{(h)}(s, a|f), \text{ for } s \in \mathbb{C}.$$

The function $H_q^{(h)}(s, a|f)$ is a meromorphic for $s \in \mathbb{C}$ with simple pole at $s = 1$, having residue $\frac{q^{ha} \log q}{q^{hf} - 1}$, and it interpolates the values

$$(14) \quad H_q^{(h)}(1-n, a|f) = -\frac{f^{n-1}}{n} q^{ha} B_{n,q^f}\left(\frac{a}{f}\right), \text{ where } n \in \mathbb{Z}, n \geq 1.$$

Now, we modify the q -analogue of the partial zeta function as follows:

$$(15) \quad H_q^{(h)}(s, a|f) = \frac{q^{ha}}{(s-1)f} a^{1-s} \sum_{j=0}^{\infty} \binom{1-s}{j} \left(\frac{f}{a}\right)^j B_{j,q^f}^{(h)}, \text{ for } s \in \mathbb{C}.$$

From (13), (14) and (15), we can derive the below:

$$(16) \quad L_q^{(h)}(s, \chi) = \frac{1}{s-1} \frac{1}{f} \sum_{a=1}^f \chi(a) q^{ha} a^{1-s} \sum_{m=0}^{\infty} \binom{1-s}{m} \left(\frac{f}{a}\right)^m B_{m,q^f}^{(h)}.$$

It is easy to see that we can express $L_q^{(h)}(s, x|\chi)$ in the similar method. Using (15) to define $H_q^{(h)}(s, a+x|f)$ for all $a \in \mathbb{Z}$ with $0 < a < f$, $x \in \mathbb{R}$ with $0 < x < 1$, we can define

$$L_q^{(h)}(s, x|\chi) = \sum_{a=1}^f \chi(a) q^{-hx} H_q^{(h)}(s, a+x|f).$$

Let f and a be the positive integers with $0 < a < f$. Then we have

$$(17) \quad L_q^{(h)}(s, x|\chi) = \frac{1}{s-1} \frac{1}{f} \sum_{a=1}^f \chi(a) q^{ha} (a+x)^{1-s} \sum_{m=0}^{\infty} \binom{1-s}{m} \left(\frac{f}{a+x} \right)^m.$$

By (17), we see that $L_q^{(h)}(s, x|\chi)$ is an analytic for $x \in \mathbb{R}$ with $0 < x \leq 1$, $s \in \mathbb{C}$, except $s \neq 1$. Furthermore, for each $n \in \mathbb{Z}$ with $n \geq 1$, we have

$$(18) \quad L_q^{(h)}(1-n, x|\chi) = -\frac{B_{n,q,\chi}^{(h)}(x)}{n}.$$

In $\mathbb{C}$, we introduced some of the basic facts about the q -extension of Dirichlet L -function which interpolates the generalized q -Bernoulli polynomials attached to χ . The values of $L_q^{(h)}(s, x|\chi)$ at negative integers are algebraic, hence may be regarded as lying in an extension of $\mathbb{Q}_p$. We therefore look for a p -adic function which agrees with at negative integers in the next section.

3. ON A p -ADIC INTERPOLATION FUNCTION FOR THE q -EXTENSION OF THE GENERALIZED BERNOULLI POLYNOMIALS AND ITS DERIVATIVE

In this section we shall consider the p -adic analogs of the q - L -functions, $L_q^{(h)}(s, x|\chi)$, which were introduced in the previous section. Indeed this functions are the q -analogs of the p -adic interpolation functions for the generalized Bernoulli polynomials attached to χ . Let w denote the *Teichmüller* character, having conductor $f_w = p^*$. For an arbitrary character χ , we define $\chi_n = \chi w^{-n}$, where $n \in \mathbb{Z}$, in the sense of the product of characters. Throughout this section, we assume that $q \in \mathbb{C}_p$ with $|1-q|_p < p^{-\frac{1}{p-1}}$. Let $\langle a \rangle = w^{-1}(a)a = \frac{a}{w(a)}$. Then, we note that $\langle a \rangle \equiv 1 \pmod{p^* \mathbb{Z}_p}$. By the definition of $\langle a \rangle$, we easily see that $\langle a+p^*t \rangle = w^{-1}(a+p^*t)(a+p^*t) = w^{-1}(a)a + w^{-1}(a)(p^*t) \equiv 1 \pmod{p^* \mathbb{Z}_p[t]}$, where $t \in \mathbb{C}_p$ with $|t|_p \leq 1$, $(a, p) = 1$. The p -adic logarithm function, $\log_p$, is the unique function $\mathbb{C}_p^\times \rightarrow \mathbb{C}_p$ that satisfy (1) $\log_p(1+x) = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} x^n$, $|x|_p < 1$, (2) $\log_p(xy) = \log_p x + \log_p y$, $\forall x, y \in \mathbb{C}_p^\times$, and $\log_p p = 0$.

Let $A_j(x) = \sum_{n=0}^{\infty} a_{n,j} x^n$, $a_{n,j} \in \mathbb{C}$, $j = 0, 1, 2, \dots$ be a sequence of power series, each of which converges in a fixed subset $D = \{s \in \mathbb{C}_p \mid |s|_p \leq |p^*|^{-1} p^{-\frac{1}{p-1}}\}$ of $\mathbb{C}_p$ such that (1) $a_{n,j} \rightarrow a_{n,0}$ as $j \rightarrow \infty$ for $\forall n$; (2) for each $s \in D$ and $\epsilon > 0$, there exists $n_0 = n_0(s, \epsilon)$ such that $\left| \sum_{n \geq n_0} a_{n,j} s^n \right|_p < \epsilon$ for $\forall j$. Then $\lim_{j \rightarrow \infty} A_j(s) = A_0(s)$ for all $s \in D$. This is used by Washington [24] to show that each of the function $w^{-s}(a)a^s$ and $\sum_{m=0}^{\infty} \binom{s}{m} \left(\frac{F}{a}\right)^m B_m$, where F is the multiple of p^* and $f = f_\chi$, is analytic in D . Let F be a positive integral multiple of p^* and $f = f_\chi$, and let

$$(19) \quad L_{p,q}^{(h)}(s, t|\chi) = \frac{1}{s-1} \frac{1}{F} \sum_{\substack{a=1 \\ (a,p)=1}}^F \chi(a) q^{ha} <a + p^*t>^{1-s} \sum_{m=0}^{\infty} \binom{1-s}{m} \left(\frac{F}{a + p^*t}\right)^m B_{m,q^F}^{(h)}.$$

Then $L_{p,q}^{(h)}(s, t|\chi)$ is analytic for $t \in \mathbb{C}_p$ with $|t|_p \leq 1$, provided $s \in D$, except $s \neq 1$ when $\chi \neq 1$. For $t \in \mathbb{C}_p$ with $|t|_p \leq 1$, we see that $\sum_{j=0}^{\infty} \binom{s}{j} \left(\frac{F}{a + p^*t}\right)^j B_{j,q^F}^{(h)}$ is analytic for $s \in D$. It readily follows that $<a + p^*t>^s = <a>^s \sum_{m=0}^{\infty} \binom{s}{m} (a^{-1} p^*t)^m$ is analytic for $t \in \mathbb{C}_p$ with $|t|_p \leq 1$ when $s \in D$. Thus, since $(s-1)L_{p,q}^{(h)}(s, t|\chi)$ is a finite sum of products of these two functions, it must also be analytic for $t \in \mathbb{C}_p$, $|t|_p \leq 1$, whenever $s \in D$. Note that

$$(20) \quad \lim_{s \rightarrow 1} (s-1)L_{p,q}^{(h)}(s, t|\chi) = \frac{1}{F} \sum_{\substack{a=1 \\ (a,p)=1}}^F \chi(a) q^{ha} B_{0,q^F}^{(h)}.$$

We now let $n \in \mathbb{Z}$, $n \geq 1$, and fix $t \in \mathbb{C}_p$ with $|t|_p \leq 1$. Since F must be a multiple of $f = f_{\chi_n}$. By (4), we see that

$$(21) \quad B_{n,q,\chi_n}^{(h)}(p^*t) = F^{n-1} \sum_{a=0}^{F-1} \chi_n(a) q^{ha} B_{n,q^F}^{(h)}\left(\frac{a + p^*t}{F}\right).$$

If $\chi_n(p) = 0$, then $(p, f_{\chi_n}) = 1$, so that $\frac{F}{p}$ is a multiple of f_{χ_n} . Therefore, we obtain

$$(22) \quad \chi_n(p) p^{n-1} B_{n,q^p,\chi_n}^{(h)}(p^{-1} p^*t) = F^{n-1} \sum_{\substack{a=0 \\ p|a}}^F \chi_n(a) q^{ha} B_{n,q^F}^{(h)}\left(\frac{a + p^*t}{F}\right).$$

The difference of these quantities yields

$$(23) \quad B_{n,q,\chi_n}^{(h)}(p^*t) - \chi_n(p)p^{n-1}B_{n,q^p,\chi_n}^{(h)}(p^{-1}p^*t) = F^{n-1} \sum_{\substack{a=1 \\ p \nmid a}}^F \chi_n(a)q^{ha}B_{n,q^F}^{(h)}\left(\frac{a+p^*t}{F}\right).$$

From the definition of q -Bernoulli polynomials (see[18:p.3]), we note that

$$\begin{aligned} B_{n,q^F}^{(h)}\left(\frac{a+p^*t}{F}\right) &= \sum_{k=0}^n \binom{n}{k} \left(\frac{a+p^*t}{F}\right)^{n-k} B_{k,q^F}^{(h)} \\ &= F^{-n}(a+p^*t)^n \sum_{k=0}^n \binom{n}{k} \left(\frac{F}{a+p^*t}\right)^k B_{k,q^F}^{(h)}. \end{aligned}$$

Since $\chi_n(a) = \chi(a)w^{-n}(a)$ and for $(a,p) = 1$, and $t \in \mathbb{C}_p$ with $|t|_p \leq 1$, we have

$$(24) \quad \begin{aligned} &B_{n,q,\chi_n}^{(h)}(p^*t) - \chi_n(p)p^{n-1}B_{n,q^p,\chi_n}^{(h)}(p^{-1}p^*t) \\ &= \frac{1}{F} \sum_{\substack{a=1 \\ (a,p)=1}}^F \chi(a)q^{ha} \langle a+p^*t \rangle^n \sum_{m=0}^{\infty} \binom{n}{m} \left(\frac{F}{a+p^*t}\right)^m B_{m,q^F}^{(h)} \end{aligned}$$

Thus, we see that

$$-\frac{1}{n} \left(B_{n,q,\chi_n}^{(h)}(p^*t) - \chi_n(p)p^{n-1}B_{n,q^p,\chi_n}^{(h)}(p^{-1}p^*t) \right) = L_{p,q}^{(h)}(1-n, t|\chi), \text{ for } n \in \mathbb{N}.$$

Therefore we obtain the following theorem:

Theorem 3. *Let F be a positive integral multiple of p^* and $f = f_\chi$, and let*

$$\begin{aligned} &L_{p,q}^{(h)}(s, t|\chi) \\ &= \frac{1}{s-1} \frac{1}{F} \sum_{\substack{a=1 \\ (a,p)=1}}^F \chi(a)q^{ha} \langle a+p^*t \rangle^{1-s} \sum_{m=0}^{\infty} \binom{1-s}{m} \left(\frac{F}{a+p^*t}\right)^m B_{m,q^F}^{(h)}. \end{aligned}$$

Then, $L_{p,q}^{(h)}(s, t|\chi)$ is analytic for $t \in \mathbb{C}_p$, $|t|_p \leq 1$, $h \in \mathbb{Z}$, provided $s \in D$, except $s \neq 1$. Also, if $t \in \mathbb{C}_p$, $|t|_p \leq 1$, this function is analytic for $s \in D$ when $\chi \neq 1$, and meromorphic for $s \in D$, with simple pole at $s = 1$ having residue $\frac{h \log_p q}{q^{hF}-1} \left(\frac{1-q^{hF}}{1-q^h} - \frac{1-q^{hF}}{1-q^{hp}} \right)$ when $\chi = 1$. Furthermore, for each $n \in \mathbb{Z}$, $n \geq 1$, we have

$$L_{p,q}^{(h)}(1-n, t|\chi) = -\frac{1}{n} \left(B_{n,q,\chi_n}^{(h)}(p^*t) - \chi_n(p)p^{n-1}B_{n,q^p,\chi_n}^{(h)}(p^{-1}p^*t) \right).$$

Remark. Note that $\lim_{h \rightarrow 0} L_{p,q}^{(h)}(s, 0|\chi) = \lim_{q \rightarrow 1} L_{p,q}^{(h)}(s, 0|\chi) = L_p(s, \chi)$ for $s \in D$ with $s \neq 1$ if $\chi = 1$, where $L_p(s, \chi)$ is Kubota-Leopoldt's p -adic L -function, cf. [5, 6, 7, 8, 11, 13, 20, 23, 24].

We now consider the q -analogue of the partial p -adic zeta function as follows:

$$H_{p,q}^{(h)}(s, a|F) = \frac{1}{s-1} \frac{1}{F} \langle a \rangle^{1-s} \sum_{j=0}^{\infty} \binom{1-s}{j} \left(\frac{F}{a}\right)^j B_{j,q^F}^{(h)},$$

where $s \in D$, $s \neq 1$, $a \in \mathbb{Z}$ with $(a, p) = 1$, and F is a multiple of p^* , cf. [8, 13]. The function $L_{p,q}^{(h)}(s, \chi)$ can be rewritten as the sum

$$L_{p,q}^{(h)}(s, \chi) = \sum_{\substack{a=1 \\ (a,p)=1}}^F \chi(a) q^{ha} H_{p,q}^{(h)}(s, a|F), \text{ cf. [8, 13],}$$

provided F is a multiple of both p^* and $f = f_\chi$. The function $H_{p,q}^{(h)}(s, a|F)$ is a meromorphic for $s \in D$ with a simple pole at $s = 1$, having residue $\frac{h \log_p q}{q^{hF} - 1}$, and it interpolates the values

$$H_{p,q}^{(h)}(1-n, a|F) = -\frac{1}{n} F^{n-1} w^{-n}(a) B_{n,q^F}^{(h)}\left(\frac{a}{F}\right),$$

where $n \in \mathbb{Z}$, $n \geq 1$, cf. [8-19]. By using $H_{p,q}^{(h)}(s, a + p^*t|F)$, we can express $L_{p,q}^{(h)}(s, t|\chi)$ for all $a \in \mathbb{Z}$, $(a, p) = 1$, and $t \in \mathbb{C}_p$ with $|t|_p \leq 1$, as follows:

$$L_{p,q}^{(h)}(s, t|\chi) = \sum_{\substack{a=1 \\ (a,p)=1}}^F \chi(a) q^{ha} H_{p,q}^{(h)}(s, a + p^*t|F).$$

It is easy to see that $H_{p,q}^{(h)}(s, a + p^*t|F)$ is analytic for $t \in \mathbb{C}_p$, $|t|_p \leq 1$, where $s \in D$, $s \neq 1$, and meromorphic for $s \in D$, with a simple pole at $s = 1$, when $t \in \mathbb{C}_p$, $|t|_p \leq 1$. Let us consider the first partial derivative of the function $L_{p,q}^{(h)}(s, t|\chi)$ at $s = 0$. The value of $\frac{\partial}{\partial s} L_{p,q}^{(h)}(0, t|\chi)$ is the coefficient of s in the expansion of $L_{p,q}^{(h)}(s, t|\chi)$ at $s = 0$. By using Taylor expansion at $s = 0$, we see that

$$\begin{aligned} \frac{1}{1-s} &= 1 + s + \cdots, \\ \langle a + p^*t \rangle^{1-s} &= \langle a + p^*t \rangle \left(1 - s \log_p \langle a + p^*t \rangle + \cdots\right), \\ \binom{1-s}{m} &= \frac{(-1)^{m+1}}{m(m-1)} s + \cdots. \end{aligned}$$

By employing these expansion, along with some algebraic manipulation, we evaluate $\frac{\partial}{\partial s} L_{p,q}^{(h)}(0, t|\chi)$. From the definition of $L_{p,q}^{(h)}(s, t|\chi)$, we note that

$$\begin{aligned} & L_{p,q}^{(h)}(s, t|\chi) \\ &= \frac{1}{s-1} \frac{1}{F} \sum_{\substack{a=1 \\ (a,p)=1}}^F \chi(a) q^{ha} <a + p^*t>^{1-s} \sum_{m=0}^{\infty} \binom{1-s}{m} \left(\frac{F}{a + p^*t} \right)^m B_{m,q^F}^{(h)}. \end{aligned}$$

Thus, we have

$$\begin{aligned} (25) \quad & \frac{\partial}{\partial s} L_{p,q}^{(h)}(s, t|\chi)|_{s=0} = \sum_{a=1}^F \chi_1(a) q^{ha} \left\{ \left(\left(\frac{a + p^*t}{F} \right) B_{0,q^F}^{(h)} + B_{1,q^F}^{(h)} \right) \log_p <a + p^*t> \right. \\ & \left. - \left(\frac{a + p^*t}{F} \right) B_{0,q^F}^{(h)} + \sum_{m=2}^{\infty} \frac{(-1)^m}{m(m-1)} B_{m,q^F}^{(h)} \left(\frac{a + p^*t}{F} \right)^{1-m} \right\} - \sum_{\substack{a=1 \\ (a,p)=1}}^F \chi_1(a) q^{ha} B_{1,q^F}^{(h)}. \end{aligned}$$

Since the Diamond gamma function is defined by

$$(26) \quad G_p(x) = (x - \frac{1}{2}) \log_p x - x + \sum_{j=2}^{\infty} \frac{B_j}{j(j-1)} x^{1-j}, \text{ for } |x|_p > 1, \text{ cf. [5, 6, 20, 23] ,}$$

and $w(a)$ is a root of unity for $(a, p) = 1$, we see that

$$\log_p <a + p^*t> = \log_p(a + p^*t) + \log_p w^{-1}(a) = \log_p(a + p^*t), \text{ cf. [24].}$$

For $f \in UD(\mathbb{Z}_p, \mathbb{C}_p) = \{f|f : \mathbb{Z}_p \rightarrow \mathbb{C}_p \text{ is uniformly differentiable function} \}$, the Volkenborn integral is defined by

$$(27) \quad I_0(f) = \int_{\mathbb{Z}_p} f(x) d\mu_0(x) = \lim_{N \rightarrow \infty} \frac{1}{p^N} \sum_{x=0}^{p^N-1} f(x), \text{ cf. [18].}$$

It is easy to see that

$$I_0(f_1) = I_0(f) + f'(0), \text{ where } f_1(x) = f(x+1).$$

If we take $f(x) = q^{hx} e^{tx}$, ($h \in \mathbb{Z}$), then we have

$$\int_{\mathbb{Z}_p} q^{hx} e^{tx} d\mu_0(x) = \frac{h \log_p q + t}{q^h e^t - 1}.$$

Thus, we note that

$$(28) \quad \int_{\mathbb{Z}_p} q^{hx} x^n d\mu_0(x) = B_{n,q}^{(h)}, \text{ for } h \in \mathbb{Z}.$$

We now consider a locally analytic function $G_{p,q}^{(h)}(x)$ which are the q -extension of Diamond gamma function as follows:

$$(29) \quad G_{p,q}^{(h)}(x) = \int_{\mathbb{Z}_p} \{(x+z) \log_p(x+z) - (x+z)\} q^{hz} d\mu_0(z), \text{ for } |x|_p > 1.$$

From the above Eq.(29), we note that $G_{p,q}^{(h)}(x)$ is locally analytic on $\mathbb{C}_p \setminus \mathbb{Z}_p$. By (28) and (29), we easily see that

$$(30) \quad G_{p,q}^{(h)}(x) = \left(x B_{0,q}^{(h)} + B_{1,q}^{(h)} \right) \log_p x - x B_{0,q}^{(h)} + \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n(n+1)} B_{n+1,q}^{(h)} x^{-n}, \text{ for } |x|_p > 1.$$

Note that $\lim_{q \rightarrow 1} G_{p,q}^{(h)}(x) = \lim_{h \rightarrow 0} G_{p,q}^{(h)}(x) = G_p(x)$. From (25) and (30), we can derive the below formula:

$$\begin{aligned} \frac{\partial}{\partial s} L_{p,q}^{(h)}(0, t|\chi) &= \sum_{\substack{a=1 \\ (a,p)=1}}^F \chi_1(a) q^{ha} G_{p,q^F}^{(h)} \left(\frac{a + p^* t}{F} \right) - L_{p,q}^{(h)}(0, \chi) \log_p F \\ &\quad - \sum_{\substack{a=1 \\ (a,p)=1}}^F \chi_1(a) q^{ha} B_{1,q^F}^{(h)}. \end{aligned}$$

Therefore we obtain the following theorem:

Theorem 4. *Let χ be the primitive Dirichlet character, and let F be a positive integral multiple of p^* and $f = f_\chi$. Then for any $t \in \mathbb{C}_p$ with $|t|_p \leq 1$, $h \in \mathbb{Z}$, we have*

$$\begin{aligned} \frac{\partial}{\partial s} L_{p,q}^{(h)}(0, t|\chi) &= \sum_{\substack{a=1 \\ (a,p)=1}}^F \chi_1(a) q^{ha} G_{p,q^F}^{(h)} \left(\frac{a + p^* t}{F} \right) - L_{p,q}^{(h)}(0, \chi) \log_p F \\ &\quad - \sum_{\substack{a=1 \\ (a,p)=1}}^F \chi_1(a) q^{ha} B_{1,q^F}^{(h)}. \end{aligned}$$

Note that

$$\lim_{h \rightarrow 0} \frac{\partial}{\partial s} L_{p,q}^{(h)}(0, t|\chi) = \frac{\partial}{\partial s} L_p(0, t|\chi) = \sum_{\substack{a=1 \\ (a,p)=1}}^F \chi_1(a) G_p\left(\frac{a+p^*t}{F}\right) - L_p(0, \chi) \log_p F.$$

This formula can be considered as the generalization of Ferrero-Greenberg's theorem, cf.[5, 6, 7, 8, 11, 13, 20, 23, 24].

REFERENCES

1. L. Carlitz, *q-Bernoulli numbers and polynomials*, Duke Math. J. **15** (1948), 987-1000.
2. M. Cenkci, M. Can, V. Kurt, *p-adic interpolation functions and Kummer type congruences for q-twisted and q-generalized twisted Euler numbers*, Advan. Stud. Contemp. Math. **9** (2004), 203-216.
3. G. Choi, A. Zaharescu, *Numerical polynomials in several variables with few nonzero terms*, Advan. Stud. Contemp. Math. **10** (2005), 7-13.
4. H.-S. Cho, E.-S. Kim, *Translation-invariant p-adic integral on $\mathbb{Z}_p$* , Notes on Number Theory and Discrete Mathematics **7** (2001), 70-77.
5. J. Diamond, *The p-adic log gamma function and p-adic Euler constant*, Trans. Amer. Math. Soc. **233** (1977), 321-337.
6. B. Ferrero, R. Greenberg, *On the behavior of p-adic L-functions at $s = 0$* , Invent. Math. **50** (1978), 91-102.
7. K. Iwasawa, *Lectures on p-adic L-functions*, Princeton Univ. Press, 1972.
8. T. Kim, *On explicit formulas of p-adic q-L-functions*, Kyushu J. Math. **48** (1994), 73-86.
9. T. Kim, *Sums powers of consecutive q-integers*, Advan. Stud. Contemp. Math. **9** (2004), 15-18.
10. T. Kim, *q-Volkenborn integration*, Russ. J. Math. Phys. **9** (2002), 288-299.
11. T. Kim, *An invariant p-adic integral associated with Daehee numbers*, Integral Trans. Special Funct. **13** (2002), 65-69.
12. T. Kim, L. C. Jang, S. H. Rim, *An extension of q-zeta function*, Int. J. Math. Math. Sci. **2004** no. **49** (2004), 2649-2651.
13. T. Kim, *On p-adic q-L-functions and sums of powers*, Discrete Math. **252** (2002), 179-187.
14. T. Kim, *On the twisted q-zeta functions and q-Bernoulli polynomials*, Far East J. Appl. Math. **13** (2003), 13-21.

15. T. Kim, *p-adic q-integrals associated with Changhee-Barnes' q-Bernoulli polynomials*, Integral Trans. Special Funct. **15** (2004), 415-420.
16. T. Kim, *Analytic continuation of multiple q-zeta functions and their values at negative integers*, Russian J. Math. Phys. **11** (2004), 71-76.
17. T. Kim, *Non-archimedean q-integrals associated with multiple Changhee q-Bernoulli polynomials*, Russian J. Math. Phys. **10** (2003), 91-98.
18. T. Kim, *A new approach to q-zeta function*, arXiv:math.NT/0502005 (1 Feb 2005).
19. T. Kim, C. Adiga, *Sums Products of generalized Bernoulli numbers*, Inter. Math. J. **5 no. 1** (2004), 1-7.
20. N. Koblitz, *A new proof of certain formulas for p-adic L-function*, Duke Math. J **46** (1979), 455-468.
21. A. A. Panchishkin, *Two variable p-adic L-functions attached to eigenfamilies of positive slope*, Invent. Math. **154** (2003), 551-615.
22. M. Schlosser, *q-analogues of the sums of consecutive integers, squares, cubes, quarts, and quints*, The Electronic J. Combinatorics **11** (2004), R 71.
23. K. Shiratani, S. Yamamoto, *On a p-adic interpolation function for the Euler numbers and its derivatives*, Mem. Fac. Sci.Kyushu Univ. **39** (1985), 113-125.
24. L. C. Washington, *Introduction to cyclotomic fields*, Springer-Verlag(1'st Ed.), 1982.